

*TNF-α* -857 and -308 ( $\chi^2 = 23.88$ ,  $P < 0.001$ ) or -238 ( $\chi^2 = 13.91$ ,  $P < 0.005$ ) and between *TNF-α* -376 and -238 ( $\chi^2 = 363.47$ ,  $P < 0.001$ ). *TNF-α* -857 T/T genotype was more frequent in patients with duodenal ulcer than in those with noncardia gastric cancer or other benign diseases, considering the patients overall ( $\chi^2 = 38.78$ ,  $P < 0.01$ ) or only those *H. pylori* infected ( $\chi^2 = 34.65$ ,  $P < 0.05$ ) or infected by *cagA* positive strains ( $\chi^2 = 12.50$ ,  $P < 0.05$ ) (Figure 1). The risk of duodenal ulcer for the *TNF-α* -857 T/T genotype was 4.28 (95% CI, 1.80–10.21). *TNF-α* -857 T allele is associated with high transcriptional activity of the promoter/enhancer region.<sup>6</sup> This might cause an enhanced mucosal release of *TNF-α* in response to *H. pylori* infection, and high mucosal levels of *TNF-α* has been suggested to play a pivotal role in the pathogenesis of duodenal ulcer.<sup>7</sup> *TNF-α* -308 SNP was correlated with *H. pylori* infection ( $\chi^2 = 9.70$ ,  $P < 0.05$ ), which was more frequent in *TNF-α* -308 A/G (61.2%) than in G/G (54.1%) subjects, in agreement with previous data by Yea et al.<sup>8</sup> in Korean patients. Antral inflammation was more severe in patients bearing *TNF-α* -1031 T allele ( $\chi^2 = 12.3$ ,  $P < 0.05$ ) (Table 1). The presence or absence of intestinal metaplasia was not correlated with any of the studied cytokines' SNPs, but it was correlated with *H. pylori* antral density grade ( $\chi^2 = 10.59$ ,  $P < 0.05$ ) and with *cagA* (Fisher exact test,  $P < 0.001$ ) or *s1 vacA* (Fisher exact test,  $P < 0.001$ ). We suggest that *TNF-α* -308 and -1031 SNPs might be involved in favoring *H. pylori* infection and the inflammatory response of the infected gastric mucosa. *TNF-α* -857 T/T genotype might predispose to duodenal ulcer. *TNF-α* SNPs in the promoter/enhancer region of the gene, do not seem involved in favoring the precancerous intestinal metaplasia or noncardia gastric cancer.

CARLO-FEDERICO ZAMBON

DANIELA BASSO

FILIPPO NAVAGLIA

ALESSANDRA FALDA

CLAUDIO BELLUCO

PAOLA FOGAR

ELIANA GRECO

NICOLETTA GALLO

FABIO FARINATI

ROMILDA CARDIN

MASSIMO RUGGE

FRANCESCO DI MARIO

MARIO PLEBANI

Departments of Medical and Surgical Sciences, Laboratory

Medicine, Oncological and Surgical Sciences,

Gastroenterology, and Pathology

University of Padova

Padova, Italy

Department of Gastroenterology

University of Parma

Parma, Italy

1. El-Omar EM, Rabkin CS, Gammon MD, Vaughan TL, Risch HA, Schoenberg JB, Stanford JL, Mayne ST, Goedert J, Blot WJ, Fraumeni JF Jr, Chow W-H. Increased risk of noncardia gastric cancer associated with proinflammatory cytokine gene polymorphisms. *Gastroenterology* 2003;124:1193–1201.
2. Basso D, Navaglia F, Brigato L, Piva MG, Toma A, Greco E, DiMario F, Galeotti F, Roveroni G, Corsini A, Plebani M. Analysis of *Helicobacter pylori* *vacA* and *cagA* genotypes and serum antibody profile in benign and malignant gastroduodenal diseases. *Gut* 1998;43:182–186.
3. Zambon C-F, Basso D, Navaglia F, Germano G, Gallo N, Milazzo M, Greco E, Fogar P, Mazza S, Di Mario F, Basso G, Rugge M, Plebani

M. *Helicobacter pylori* virulence genes and host IL-1RN and IL-1β genes interplay in favouring the development of peptic ulcer and intestinal metaplasia. *Cytokine* 2002;18:242–251.

4. Montecucco C, Rappuoli R. Living dangerously: how *Helicobacter pylori* survives in the human stomach *Nature Rev Mol Cell Biol* 2001;2:457–466.
5. El-Omar EM, Carrington M, Chow W-H, McColl KEL, Bream JH, Young HA, Herrera J, Lissowska J, Yuan C-C, Rothman N, Lanyon G, Martin M, Fraumeni JF Jr, Rabkin CS. Interleukin-1 polymorphisms associated with increased risk of gastric cancer. *Nature* 2000;404:398–402.
6. Tsukasaki K, Miller CW, Kubota T, Takeuchi S, Fujimoto T, Ikeda S, Tomonaga M, Koeffler HP. Tumor necrosis factor alpha polymorphism associated with increased susceptibility to development of adult T-cell leukemia/lymphoma in human T-lymphotropic virus type 1 carriers. *Cancer Res* 2001;61:3770–3774.
7. Alkout AM, Blackwell CC, Weir DM. Increased inflammatory responses of persons of blood group O to *Helicobacter pylori*. *J Infect Dis* 2000;181:1364–1369.
8. Yea SS, Yang Y-I, Jang WH, Lee YJ, Bae H-S, Paik K-H. Association between *TNF-α* promoter polymorphism and *Helicobacter pylori* *cagA* subtype infection. *J Clin Pathol* 2001;54:703–706.

doi:10.1053/j.gastro.2003.08.042

**Reply.** We thank Zambon et al. for their interest in our paper and for sharing their data. Their study, comprising a heterogeneous group of upper gastrointestinal pathologies and including 129 gastric cancer cases, failed to find statistically significant associations between *IL-1B*-31 and *TNFA*-308 proinflammatory genotypes and gastric cancer or its precursors. However, their findings are at odds with a number of larger studies that have looked at these important markers. For example, there were 366 gastric cancer cases vs. 429 population controls in our earlier study based in Poland,<sup>1</sup> and this was confirmed in our U.S. study with 188 cases and 210 controls.<sup>2</sup> Machado et al. independently confirmed both *IL-1B*-511<sup>3,4</sup> and *TNFA*-308 findings in a study of 287 gastric cancer cases vs. 306 controls.<sup>5</sup> Furthermore, Rad et al.<sup>6</sup> and Furuta et al.<sup>7</sup> confirmed the association between *IL-1* markers and gastric premalignant abnormalities in Caucasian and Japanese populations, respectively.

We found it difficult to analyze or comment on the other findings in the study by Zambon et al. due to the lack of subject details and restriction of statistical data to  $\chi^2$  and  $P$  values. We also caution about their reported associations with the *TNFA* SNP's -1031 and -857 pending better definition of the functional significance of these genetic markers. Nevertheless, we agree with Zambon et al. about the important role of cytokine gene polymorphisms in the pathogenesis of *H. pylori*-related diseases.

EMAD M. EL-OMAR

Department of Medicine and Therapeutics

Aberdeen University

Aberdeen, Scotland

CHARLES S. RABKIN

WONG-HO CHOW

Division of Cancer Epidemiology and Genetics

National Cancer Institute

Bethesda, Maryland

1. El-Omar EM, Carrington M, Chow WH, McColl KE, Bream JH, Young HA, Herrera J, Lissowska J, Yuan CC, Rothman N, Lanyon G, Martin M, Fraumeni JF Jr, Rabkin CS. Interleukin-1 polymorphisms associated with increased risk of gastric cancer. *Nature* 2000;404:398–402.
2. El-Omar EM, Rabkin CS, Gammon MD, Vaughan TL, Risch HA,

- Schoenberg JB, Stanford JL, Mayne ST, Goedert J, Blot WJ, Fraumeni JF Jr, Chow WH. Increased risk of noncardia gastric cancer associated with proinflammatory cytokine gene polymorphisms. *Gastroenterology* 2003;124:1193–1201.
3. Machado JC, Pharoah P, Sousa S, Carvalho R, Oliveira C, Figueiredo C, Amorim A, Seruca R, Caldas C, Carneiro F, Sobrinho-Simoes M. Interleukin 1B and interleukin 1RN polymorphisms are associated with increased risk of gastric carcinoma. *Gastroenterology* 2001;121:823–829.
  4. Figueiredo C, Machado JC, Pharoah P, Seruca R, Sousa S, Carvalho R, Capelinha AF, Quint W, Caldas C, Van Doorn LJ, Carneiro F, Sobrinho-Simoes M. *Helicobacter pylori* and interleukin 1 genotyping: an opportunity to identify high-risk individuals for gastric carcinoma. *J Natl Cancer Inst* 2002;94:1680–1687.
  5. Machado JC, Figueiredo C, Canedo P, Pharoah P, Carvalho R, Nabais S, Castro AC, Campos ML, Van Doorn LJ, Caldas C, Seruca R, Carneiro F, Sobrinho-Simoes M. A proinflammatory genetic profile increases the risk for chronic atrophic gastritis and gastric carcinoma. *Gastroenterology* 2003;125:364–371.
  6. Rad R, Prinz C, Neu B, Neuhofer M, Zeitner M, Volland P, Becker I, Schepp W, Gerhard M. Synergistic effect of *Helicobacter pylori* virulence factors and interleukin-1 polymorphisms for the development of severe histological changes in the gastric mucosa. *J Infect Dis* 2003;188:272–281.
  7. Furuta T, El Omar EM, Xiao F, Shirai N, Takashima M, Sugimura H. Interleukin 1beta polymorphisms increase risk of hypochlorhydria and atrophic gastritis and reduce risk of duodenal ulcer recurrence in Japan. *Gastroenterology* 2002;123:92–105.

---

doi:10.1053/j.gastro.2003.11.033

## Discussion on Impaired Expression of Peroxisome Proliferator-Activated Receptor $\gamma$ in Ulcerative Colitis

Dear Sir:

In their interesting paper on peroxisome proliferator-activated receptor (PPAR $\gamma$ ) expression in inflammatory bowel disease (IBD), Dubuquoy et al.<sup>1</sup> examined colonic biopsies from patients with ulcerative colitis (UC), and Crohn's disease (CD). For control groups they used biopsies from patients with irritable bowel syndrome and sigmoid diverticulitis. We feel that clarification is required as to the true nature of the latter group in relation to its use as an inflammatory control.

Diverticulitis is a subserosal disease.<sup>2</sup> Narrow-necked diverticula can become obstructed by entrapped fecal matter. Hyperplasia of the

mucosa-associated lymphoid tissue follows. Subsequent inflammation of the pericolic and mesenteric fat can lead to peridiverticular abscess formation. Mucosal inflammation, however, is not a prominent feature of diverticulitis.

In contrast, diverticular colitis is characterized by segmental mucosal inflammation, the proposed etiology of which is multifactorial.<sup>3</sup> Potential mechanisms include the mechanical action of firm stools on redundant mucosal folds leading to mucosal ischemia and prolonged exposure of the mucosa to luminal contents such as bacterial antigens and toxins. Endoscopically, mucosal hyperemia, granularity, edema, and ulceration can be identified on crescentic mucosal folds; diverticular orifices are usually spared.<sup>4</sup> The microscopic features include crypt abscesses and architectural distortion and are often indistinguishable from those found in idiopathic IBD.<sup>5</sup>

Since diverticular colitis is a predominantly mucosal disease, patients with this diagnosis would seem to be a more appropriate inflammatory control group than those with diverticulitis, which is primarily subserosal. An alternative and perhaps more readily accessible inflammatory control group would of course be patients with infective colitis.

RICHARD MAKINS, B.Sc., M.R.C.P.

DAVID RAMPTON, DPHIL, F.R.C.P.

*Academic Department Adult & Paediatric Gastroenterology*

*Baits and The London*

*Queen Mary's School of Medicine and Dentistry*

*London, England*

1. Dubuquoy L, Jansson EA, Deeb S, Rakotobe S, Karoui M, Colombel J-F, Auwerx J, Pettersson S, Desreumaux P. Impaired expression of peroxisome proliferator activated receptor  $\gamma$  in ulcerative colitis. *Gastroenterology* 2003;124:1265–1276.
2. Ludeman L, Warren BF, Shepherd NA. The pathology of diverticular disease. *Best Pract Res Clin Gastroenterol* 2002;16:543–562.
3. Ludeman L, Shepherd NA. What is diverticular colitis? *Pathology* 2002;34:568–572.
4. Gore S, Shepherd NA, Wilkinson SP. Endoscopic crescentic fold disease of the sigmoid colon: the clinical and histopathological spectrum of a distinctive endoscopic appearance. *Int J Colorectal Dis* 1992;7:76–81.
5. Shepherd NA. Pathological mimics of chronic inflammatory bowel disease. *J Clin Pathol* 1991;44:726–733.

---

doi:10.1053/j.gastro.2003.08.043